

A possible connection between gamma-ray bursts and single black holes

G.M. Beskin¹, A. Shearer², A. Golden², C. Bartolini³, A. Guarnieri³, A. Piccioni³, and N. Masetti⁴

¹ Special Astrophysical Observatory, Russia

² University of Galway, Ireland

³ Dipartimento di Astronomia, Università di Bologna, via Zamboni 33, I-40126 Bologna, Italy

⁴ Istituto Te.S.R.E., CNR, Bologna, Italy

Received December 18, 1998; accepted July 1, 1999

Abstract. We consider Single Black Holes (SBHs) of $10^3 M_{\odot}$ as possible parent objects for cosmological GRBs. The latter are the result of the energy release of SBHs magnetosphere inhomogeneities. The percolation nature of this release yields a unique power law for the GRBs energy distribution that is $E^{-\beta}$ (with $\beta = 1.7 - 1.8$), i.e. the same observed in solar flares, flare stars and X-ray sources. The number of GRBs per year per galaxy is about $10^{-2} - 10^{-1}$ with total energy $10^{47} - 10^{50}$ ergs for beaming factor of about $10^{-4} - 10^{-5}$.

Key words: black hole physics, gamma-rays: bursts

1. Introduction

There is much evidence that most GRBs have cosmological origin (see Piran 1998 for a recent review). Observations give total isotropic energy fluences of these events in the range $10^{51} - 10^{54}$ erg during several seconds. From BATSE observations, the GRBs rate is one burst per million years per galaxy. The energy release and rate can be reduced, of course, by several orders of magnitude if the GRBs are beamed with a factor of about of $10^{-4} - 10^{-5}$. The central point in the GRBs scenario is the origin of the “inner engine” that we do not detect directly. A significant difficulty with many proposed parent bodies of GRBs is that only a small amount of baryons ($< 10^{-5} M_{\odot}$) can be involved if a required relativistic expansion is to be produced (Shemi & Piran 1990). Scenarios using coalescence of two neutron stars (Narayan et al. 1992) or special types of supernovae (Paczynski 1998) involve potentially large amounts of “contaminating” baryons. A baryon-poor mechanism is based on the possibility to convert rotational energy of neutron stars due to strong magnetic field (10^{15} G) to a Poynting flux and then, to kinetic energy of particles (see e.g. Usov 1992). We proposed as similar sources of magnetic energy, the inhomogeneities in the magnetospheres of Single Black Holes (SBH) of $(10 - 10^3)M_{\odot}$ (Beskin et al.

1998). These objects can be parent bodies of both galactic and cosmological GRBs.

In this paper we discuss in detail a possible connection between cosmological GRBs and SBHs.

2. Assumptions for the GRB mechanism

During accretion of interstellar plasma onto a SBH, magnetic fields strengthen up to $10^{11} - 10^{12}$ G and probably form a regular magnetosphere (Bisnovatyj-Kogan & Ruzmaikin 1976; Kardashev 1995; Chakrabarti 1996; Beskin 1997).

In the magnetosphere there are local inhomogeneities similar to those of Sun and flare stars (Byrne 1992; Priest 1994) but with magnetic fields up to 10^{14} G.

The energy of these inhomogeneities as the result of spontaneous development of plasma instabilities turns into energy of fast particles and into anomalous heating. The GRB is therefore the burst of photons that have been generated as the result of such a process. We think this is an analog of solar flares and flashes of UV Ceti type stars and of the activity in X-ray binaries with compact components (for some evidence of this see Katsova & Livshits 1991; Rotschild et al. 1974; Meekins 1984; Bartolini et al. 1994).

The distribution of these flares (as well as GRBs) as a function of energy has a universal character, i.e. $\frac{dN(E)}{dE} \propto E^{-\beta}$, where $\beta = 1.7 - 1.8$. Such a universality is caused by the percolation nature of the stability break in active regions (Feder 1988; Wentzel & Seiden 1992; Pustil'nik 1997). Electrons (or positrons) and protons are accelerated in processes of dissipation of magnetic energy inhomogeneities up to $\gamma \sim 10^3 - 10^6$. Particle streams are beamed as result of one- and two-dimensions structure of magnetic field reconnection regions. They can change the direction of the motion very rapidly. The radiation is generated by these particles in magnetic fields near SBHs and in the interstellar medium (ISM). The emission is beamed as result of the stream sharpness and of the relativistic motion.

3. Energy and number of GRBs

The SBHs under consideration are remnants of Population I-II and III stars located in the disc or in the halo of galaxies. In the first case their masses are greater than $10 M_\odot$, while in the second are greater than $10^3 M_\odot$. In this hypothesis, SBHs are made of baryonic dark matter and have integral mass about $(0.1 - 1)M_{\text{gal}} \sim (10^9 - 10^{10})M_\odot$; this means that the number of SBHs is $10^6 - 10^8$ per galaxy. These estimates come from dynamical, photometric and chemical data (Bahcall 1984; Carr 1994). Of course the density of SBHs has to be higher in galaxies with star-forming regions.

We show here that the maximum energy of cosmological GRBs can be accumulated in magnetospheric inhomogeneities near massive ($M \gtrsim 10^3 M_\odot$) SBHs. Observing isotropic energy of GRBs is $10^{51} - 10^{54}$ ergs (Piran 1998). There are many reasons to think that their radiation is beamed with a factor $\sigma = \theta^2/4\pi \sim 10^{-4} - 10^{-5}$. θ is either angular size of particles stream or beaming angle of these particles radiation ($1/\gamma$) (Piran 1998). After taking into account this effect, the energy of GRBs becomes $10^{47} - 10^{50}$ ergs in a rest frame. It was shown the SBH accretes the plasma from ISM with a rate (for recent review see Chakrabarti 1996) $\dot{M} = 1.4 \cdot 10^{13} n M_3^2 V_{300}^{-3} \text{g s}^{-1}$, where n is number density of ISM in units of cm^{-3} , M_3 is the SBH mass in units of $10^3 M_\odot$, V_{300} is velocity of SBH in units of 300 km s^{-1} . The magnetic field frozen in the plasma during accretion is reinforced and conserved near the horizon (e.g. Bisnovatj-Kogan & Ruzmaikin 1976; Macdonald et al. 1986; Kardashev 1995; Beskin 1997). There is not a self-consistent model of this process and we will estimate possible parameters for the SBH magnetosphere in a phenomenological frame only.

If the part of the full mass-energy of the accreted plasma $\delta \dot{M} c^2$ is stored in form of magnetosphere energy during the time t , then $(H_g/8\pi)(4/3)\pi r_g^3 \sim \delta \dot{M} c^2 t$, where H_g value of global magnetic field, $r_g = 2GM/c^2$, and $H_g \sim 3 \cdot 10^{12} \delta_{-1}^{1/2} M_3^{-1/2} V_{300}^{-3/2} n^{1/2} t_9^{1/2} \text{ G}$, where δ_{-1} is expressed in units of 0.1, and t in units of 10^9 years.

For a magnetic field H_1 in inhomogeneities of size r for constant magnetic flux it is easy to obtain that: $H_1 \sim H_g (r_g/r)^2 \sim 3 \cdot 10^{14} \alpha_{0.1}^{-2} \delta_{-1}^{1/2} M_3^{-1/2} V_{300}^{-3/2} n^{1/2} t_9^{1/2} \text{ G}$, where $\alpha = r_g/r$ is in units of 0.1.

For the magnetic energy concentrated in the inhomogeneities we obtain $E_1 \sim (H_1^2/8\pi) \cdot (4/3)\pi r^3 \sim 4 \cdot 10^{50} \alpha_{0.1}^{-1} M_3^3 \delta_{-1} n V_{300}^{-3} t_9$ ergs. This means that this amount of energy, accumulated during 10^9 years, can be released as observable GRBs. We estimate that the mass of matter trapped in the inhomogeneity region during the time t is

$$m = \dot{M} (r/r_g)^3 t \sim 4.3 \cdot 10^{26} n M_3^2 V_{300}^{-3} \alpha_{0.1}^3 t_9 \sim 2 \cdot 10^{-7} M_\odot n M_3^2 V_{300}^{-3} \alpha_{0.1}^3 t_9.$$

Only $2 \cdot 10^{-7} M_\odot$ is accumulated during 10^9 years. This means that the baryon contamination is absent. In the results of instabilities development in inhomogeneity its

magnetic energy converts to kinetic energy of electrons and protons. If β is the efficiency of this conversion, then $\beta n_e \gamma_e m_e c^2 \sim \beta n_p \gamma_p m_p c^2 \sim (H_1^2)/8\pi$, and we obtain $\gamma_e \sim 10^6 \beta$, $\gamma_p \sim 10^3 \beta$. The stream of particles is extremely relativistic as it is necessary to explain the GRBs observational properties.

We postulate a universal form for flares number distribution with energy if they are connected with the magnetosphere in inhomogeneities. This means that the parameter k of the distribution $dN(E)/dE = kE^{-\beta}$ ($\beta \sim 1.7$) is the same for solar and UV Ceti flares stars and GRBs, they have different energy ranges: $10^{27} - 10^{32}$ ergs for Sun and UV Ceti stars, $10^{47} - 10^{50}$ ergs for GRBs. Taking $k \sim 6 \cdot 10^{24} \text{ yr}^{-1}$ from observations of UV Ceti stars (Beskin et al. 1988) we obtain for a minimal GRBs energy of 10^{47} ergs (in the rest frame) about 10^{-8} flares per SBH per year. Using estimated number of SBH per galaxy, $10^6 - 10^8$, we have $10^{-2} - 10^0$ GRBs per year per galaxy. Piran (1998) gives for this number 10^{-2} (for beaming factor $\sigma \sim 10^{-4}$). Taking into account all of our assumptions, this coincidence is very promising!

Acknowledgements. This investigation was supported by the Russian Fund of Fundamental Research, by the Educational Scientific Centre "Cosmion", by the Italian Ministry of Foreign Affairs and by the University of Bologna (Funds for selected research topics).

References

- Bahcall J., 1984, ApJ 219, 1008
- Bartolini C., et al., 1994, ApJS 92, 455
- Beskin V.S., 1997, Uspechi Fiz. Nauk 167, 689, Gershberg R.E. (eds.). Flares and Flashes, Springer, Berlin, p. 330
- Beskin G.M., et al., 1998, in: Meegan C.A., Preece R.D., Koshut T.M. (eds.) 4th Huntsville GRB Symposium, AIP Conf. Proc. 428, 815
- Beskin G.M., et al., 1988, Bull. KrAO 79, 71
- Bisnovatj-Kogan, Ruzmaikin, 1976, Ap. Space Sci. 42, 401
- Byrne P.B., 1992, in: Linsky J.L., Serio S. (eds.) Physics of Solar & Stellar Coronae. Kluwer, Dordrecht, p. 489
- Carr B., 1994, ARA&A 32, 531
- Chakrabarti S.K., 1996, Phys. Rep. 266, 229
- Feder J., 1988, Fractals. Plenum Press, New York
- Kardashev N.S., 1995, MNRAS 276, 515
- Katsova M.M., Livshits M.A., 1991, SvA 68, 131
- Macdonald D.A., et al., 1986, in: Thorne K. et al. (eds.). Black Holes: The Membrane Paradigm, Yale Univ. Press
- Meekins J.F., 1992, ApJ 278, 288
- Narayan R., Paczynski B., Piran T., 1992, ApJ 395, L83
- Paczynski B., 1998, ApJ 494, L45
- Piran T., 1998, Phys. Rep. (in press) (astroph/9810256)
- Priest E.R., 1994, in: Greiner J., Dürbeck H.W., Gershberg R.E. (eds.). Flares and Flashes, Springer, Berlin, p. 3
- Pustil'nik L.A., 1997, Ap. Space Sci. 252, 325
- Rothschild R.E., et al., 1974, ApJ 189, L13
- Shemi A., Piran T., 1990, ApJ 365, L55
- Usov V.V., 1992, Nat 357, 472
- Wentzel D.G., Seiden P.E., 1992, ApJ 390, 280