

Erratum

BeppoSAX LECS background subtraction techniques

A.N. Parmar¹, T. Oosterbroek¹, A. Orr¹, M. Guainazzi¹, N. Shane^{1,2}, M.J. Freyberg³, D. Ricci⁴, and A. Malizia^{4,5}

¹ Astrophysics Division, Space Science Department of ESA, ESTEC, Postbus 299, 2200 AG Noordwijk, The Netherlands

² Pembroke College, University of Oxford, St Aldates, Oxford, OX1 1DW, UK

³ Max-Planck-Institut für Extraterrestrische Physik, D-85740 Garching bei München, Germany

⁴ BeppoSAX Science Data Center, Nuova Telespazio, via Corcolle 19 I-00131 Roma, Italy

⁵ Department of Physics and Astronomy, Southampton University, SO17 1BJ, UK

Astron. Astrophys. Suppl. Ser. **136** (1999) 407-418

Table 3 in Parmar et al. contains an error: the two first coefficients printed in this table were inadvertently exchanged. The correct table is reproduced below.

Table 3. Coefficients of the fifth-order polynomial fit to the ratio of additional counts in the background semi-annuli divided by source counts as a function of energy in keV. These coefficients can be used to correct counts extracted from the background semi-annuli for source contamination. a_0 is the constant term

a_n	Value	a_n	Value
a_0	$+4.11 \cdot 10^{-3}$	a_3	$-6.13 \cdot 10^{-5}$
a_1	$-2.71 \cdot 10^{-3}$	a_4	$-1.01 \cdot 10^{-5}$
a_2	$+9.79 \cdot 10^{-4}$	a_5	$+9.57 \cdot 10^{-7}$